

REDISCOVERED PSEUDOSCORPION TYPE MATERIAL DESCRIBED BY BEIER
FROM SOUTHEASTERN QUEENSLAND (ARACHNIDA: PSEUDOSCORPIONIDA)

MARK S. HARVEY

Harvey, M.S. 1994 12 01: Rediscovered pseudoscorpion type material described by Beier from southeastern Queensland (Arachnida: Pseudoscorpionida). *Memoirs of the Queensland Museum* 37(1):155-156. Brisbane. ISSN 0079-8835.

All or part of the type series of five pseudoscorpion species described by Beier (1969) from southeastern Queensland have been considered lost for some time. They have been recently found in the Department of Entomology, University of Queensland and transferred to the Queensland Museum. The specimens are documented. □ *Arachnida, Pseudoscorpionida, Tridenchthoniidae, Chthoniidae, type material.*

Mark S. Harvey, Western Australian Museum, Francis Street, Perth, Western Australia 6000, Australia; 15 December 1993.

Beier (1969) recorded several species of pseudoscorpions from southeastern Queensland, based upon material sent to him by Mr J.B. Williams. Five were new and a further two species were recorded from southeastern Queensland for the first time. Beier claimed to have returned most of the material to Mr Williams, then at Monash University, Melbourne, while retaining several specimens of two species in the collections of the Naturhistorisches Museum, Vienna. Despite efforts to trace these specimens, the material appeared to be lost (Harvey, 1981, 1985, 1989).

During a recent visit to the Department of Entomology, University of Queensland, Brisbane, I located the missing vials, and now take this opportunity to comment on the condition and status of the types.

All of the type material has been transferred to the Queensland Museum, Brisbane (QM).

Family TRIDENCHTHONIIDAE

Heterolophus australicus Beier, 1969

Heterolophus australicus Beier, 1969: 171-172, fig. 1.

MATERIAL EXAMINED

HOLOTYPE: ♀, Mt Tamborine, Joalah National Park, Queensland, 28°56'S, 153°12'E, berlesate, rainforest leaf litter, 16 Aug 1966, B. Cantrell, QMS21333.

PARATYPE: 1 tritonymph, same data as holotype, QMS21334.

REMARKS

The specimens are in fair condition, and the right chela is detached from the holotype. A recurrent typographical error by Beier (1969) referred to Mt Tamborine as "Mt Jamborine" and Joalah National Park as "Toalah National Park".

Anaulacodithella australica Beier, 1969

Anaulacodithella australica Beier, 1969: 172-174, fig. 2.

MATERIAL EXAMINED

HOLOTYPE: ♂ Lamington National Park, Queensland, 27°12'S, 153°10'E, extracted from leaf litter, 6 Mar 1965, B. Cantrell, QMS21335.

ALLOTYPE: ♀, Mt Glorious, Queensland, 27°20'S, 152°46'E, 11 Jul 1965, berlesate, rainforest leaf litter, B. Cantrell, QMS21336.

PARATYPES: 1 ♂, 1 ♀, Landsborough, Queensland, 26°48'S, 152°58'E, Australia, leaf mould, mixed forest, 16 Jul 1965, F.A. Perkins, QMS21337; 8 specimens, Mt Clunie, east spur, Queensland, 28°20'S, 152°31'E, leaf mould, rainforest, 5 Apr 1953, T.E. Woodward, QMS21338; 16 specimens, Mt Tamborine, Joalah National Park, Queensland, 28°56'S, 152°12'E, berlesate, 16 Aug 1966, B. Cantrell, QMS21339.

REMARKS

The holotype and allotype are in fair condition, but the remaining specimens are extremely bleached and fragmented. The left chela of the holotype has been detached; the right chela of the allotype has been detached and subsequently lost.

Family CHTHONIIDAE

Morikawia queenslandica Beier, 1969

Morikawia queenslandica Beier, 1969: 174-175, fig. 3.

MATERIAL EXAMINED

SYNTYPES: 5 ♂, 3 ♀, Mt Tamborine, Joalah National Park, Queensland, 28°56'S, 153°12'E, berlesate, rainforest leaf litter, 16 Aug 1966, B. Cantrell, QMS21341.

REMARKS

This species was transferred to the genus *Tyrannochthonius*, based on examination of the remaining syntypes lodged in Naturhistorisches Museum, Vienna (Harvey, 1989).

Morikawia semihorrida Beier, 1969

Morikawia semihorrida Beier, 1969: 176-177, fig. 4.

MATERIAL EXAMINED

HOLOTYPE: ♀, Mt Nebo, Queensland, 27°23'S, 152°47'E, rainforest leaf litter, 28 Mar 1967, J.B. Williams, QMS21340.

REMARKS

The recent transfer of this species to *Tyrannochthonius* (Harvey, 1989) based solely on the original description is confirmed by examination of the holotype. The right chela has been detached from the pedipalp.

Pseudotyrannochthonius queenslandicus
Beier, 1969

Pseudotyrannochthonius queenslandicus Beier, 1969: 177-178, fig. 5.

MATERIAL EXAMINED

HOLOTYPE: ♂, Mt Tamborine, Joalah National Park, Queensland, 28°56'S, 153°12'E, 16 Aug 1966. B. Cantrell, berlese, QMS21342.

PARATYPE: 1 tritonymph?, same data as holotype, QMS21343.

REMARKS

The specimens are in fair condition.

Austrochthonius australis Hoff, 1951

REMARKS

Beier (1969) recorded this species from southeastern Queensland, based on one male from Mt Nebo, (berlesate, rainforest leaf litter, 28 Mar 1967, J.B. Williams), which is now lodged in the Department of Entomology, University of

Queensland. This specimen conforms to recent descriptions of the species (Kennedy, 1989).

Family OLPHIDAE

Xenolpium longiventer (L. Koch, 1885)

REMARKS

Beier (1969) recorded this species from southeastern Queensland based on a specimen from Mt Nebo (rainforest leaf litter, 28 Mar 1967, J.B. Williams), which is now lodged in the Department of Entomology, University of Queensland. The specimen is clearly not congeneric with the type of *X. longiventer* (BMNH), and in fact represents an undescribed species of *Protogarypinus*. Members of this genus are abundant in southeastern Australia, and many new species await description.

ACKNOWLEDGEMENTS

Ms Margaret Schneider very kindly allowed access to the collections of the Department of Entomology, University of Queensland.

LITERATURE CITED

- BEIER, M. 1969. Neue Pseudoskorpione aus Australien. *Annalen des Naturhistorischen Museums in Wien* 73: 171-187.
- HARVEY, M.S. 1981. A checklist of the Australian Pseudoscorpionida. *Bulletin of the British Arachnological Society* 5: 237-252.
1985. Pseudoscorpionida. Pp.126-155. In Walton, D.W. (ed.), 'Zoological catalogue of Australia'. Vol. 3. (Australian Government Publishing Service: Canberra).
1989. Two new cavernicolous chthoniids from Australia, with notes on the generic placement of the south-western Pacific species attributed to the genera *Paraliochthonius* Beier and *Morikawia* Chamberlin (Pseudoscorpionida: Chthoniidae). *Bulletin of the British Arachnological Society* 8: 21-29.
- KENNEDY, C.M.A. 1989. Redescription of *Austrochthonius australis* Hoff (Chthoniidae: Pseudoscorpionida). *Proceedings of the Linnean Society of New South Wales* 111: 233-240.